

Suppressing Isocurvature Perturbations of QCD Axion Dark Matter

Kwang Sik Jeong* and Fuminobu Takahashi†

Department of Physics, Tohoku University, Sendai 980-8578, Japan

We propose a novel mechanism to suppress the isocurvature perturbations of the QCD axion. The point is that the QCD interactions become strong at an intermediate or high energy scale in the very early Universe, if the Higgs field has a sufficiently large expectation value. The effective QCD scale can be even higher in the presence of extra colored particles. We show that the QCD axion then becomes so heavy during inflation that its isocurvature perturbations are significantly suppressed, thereby relaxing the constraint on the inflation scale.

The axion a is a Nambu-Goldstone boson associated with the spontaneous breakdown of the Peccei-Quinn (PQ) symmetry introduced to solve the strong CP problem in quantum chromodynamics (QCD) [1, 2]. The axion acquires a tiny but non-zero mass from the QCD instanton effects, and it is stabilized at the CP conserving vacuum. The dynamical relaxation of the CP phase necessarily induces coherent oscillations of the axion, which contribute to cold dark matter (CDM) as the axion is stable in a cosmological time scale.

If the axion is present during inflation, it is subject to quantum fluctuations,

$$\delta a \simeq \frac{H_{\text{inf}}}{2\pi}, \quad (1)$$

where H_{inf} denotes the Hubble parameter during inflation. The quantum fluctuations lead to an almost scale-invariant CDM isocurvature density fluctuation, which leaves a distinctive imprint on the cosmic microwave background (CMB) spectrum. The observed CMB spectrum can be well fitted by a nearly scale-invariant adiabatic density perturbation, and a mixture of the isocurvature perturbation is tightly constrained. This constraint can be interpreted as an upper bound on the inflation scale in the axion CDM scenario [3],

$$H_{\text{inf}} < 0.87 \times 10^7 \text{ GeV} \left(\frac{f_a}{10^{11} \text{ GeV}} \right)^{0.408}, \quad (2)$$

at 95% CL, where f_a is the axion decay constant [4, 5]. The constraint becomes much more stringent for a smaller value of f_a due to anharmonic effects [6, 7]. Such a tight constraint on

the inflation scale can be satisfied only for small-field inflation. In particular, a chaotic inflation model [8–11] is in tension with the axion CDM.

There are several known ways to avoid the constraint. First, if the PQ symmetry is restored during inflation, the axion does not exist, and so there is no isocurvature perturbation. The axion appears when the PQ symmetry is spontaneously broken after inflation. In this case, however, topological defects such as cosmic strings and domain walls are generated, and in particular, the PQ sector must be such that the domain wall number is unity. Secondly, if the axion decay constant during inflation is larger than the present one, the isocurvature constraint can be relaxed [12].¹ This is especially the case if the bosonic partner of the axion, saxion, has a relatively flat potential and plays an important cosmological role.

In this letter we propose another way to relax the isocurvature constraint on the inflation scale in the axion CDM scenario. The point is that, if the Higgs field has a sufficiently large expectation value during inflation, the QCD confines at an intermediate or high energy scale. The QCD scale can be higher in the presence of additional colored particles that become heavy during inflation. Then, the axion becomes so heavy that its quantum fluctuations at superhorizon scales are significantly suppressed, thereby relaxing the constraint on the inflation scale.

The stronger QCD in the early Universe was considered in Refs. [14, 15] in order to suppress the cosmological abundance of the axion, which however turned out to be rather non-trivial [16].

*email: ksjeong@tuhep.phys.tohoku.ac.jp

†email: fumi@tuhep.phys.tohoku.ac.jp

¹ This has been studied in a context of non-minimal coupling to gravity in a very recent paper [13].

There are a variety of scenarios leading to a large field value of the Higgs fields [17–19]. Indeed this can naturally be realized in the supersymmetric (SUSY) standard model (SM) which possesses flat directions involving the Higgs fields H_u and H_d . In the rest of this letter we consider a SUSY axion model, but our main idea can be applied to the non-SUSY case as well as other string theoretic axions in a straightforward way.

We now examine how heavy the axion can be during inflation when the $H_u H_d$ flat direction has a large field value. In the following we assume that the Higgs fields are neutral under the PQ symmetry.² The relevant scales are the axion decay constant f_a , the effective QCD scale Λ_h , and the inflation scale H_{inf} . For simplicity we assume that the Kähler potential is a generic one such that soft scalar masses are of order the gravitino mass $m_{3/2}$, and that f_a remains constant during and after inflation.

Let ϕ^2 parameterize the $H_u H_d$ flat direction. The ϕ acquires a negative Hubble-induced mass term from the coupling with the inflaton X ,

$$\Delta K = c|X|^2|\phi|^2, \quad (3)$$

for $c > 1$, which drives ϕ to a large field value for $H_{\text{inf}} > m_{3/2}$. The coefficient c is expected to be of order unity. Here we adopt the Planck unit in which $M_{Pl} \simeq 2.4 \times 10^{18}$ GeV is set to be unity. The flat direction is stabilized at $\phi_0 \sim \sqrt{H_{\text{inf}} M}$ by balancing the negative Hubble-induced mass squared with the higher order superpotential term,

$$\Delta W = \frac{\phi^4}{M}, \quad (4)$$

where M is an effective cut-off scale. We will assume ϕ_0 to be around or above the GUT scale, $M_{\text{GUT}} \simeq 2 \times 10^{16}$ GeV, which can be realized for an appropriate value of M . It is also possible to stabilize ϕ by a higher order term in the Kähler potential. Note that the mass of the radial component of ϕ is of order H_{inf} . The phase component of ϕ can have a mass of similar size in the presence

of interactions like $X|\phi|^2$ or $|X|^2\phi^2$ in the Kähler potential. In this case there are no light degrees of freedom associated with the flat direction.³

The SSM matter fields including neutrinos acquire large SUSY mass from the Yukawa couplings with H_u or H_d . Let us first consider a case in which all the quarks are heavier than Λ_h . One can then integrate them out and obtain a pure SUSY $SU(3)_c$ with the gauge kinetic function,

$$f_h = (\text{constant}) - \frac{n}{8\pi^2} \ln S - \frac{N_f}{8\pi^2} \ln \phi, \quad (5)$$

where S denotes the axion superfield, and $N_f = 6$ is the flavor number of the SM quarks. The constant n counts the number of PQ charged quarks when the PQ symmetry is linearly realized. If Λ_h is larger than H_{inf} , the gluino condensate is formed during inflation, giving rise to the non-perturbative superpotential,

$$W_{\text{np}} = N_c \Lambda_0^3 \propto e^{-8\pi^2 f_h / N_c}, \quad (6)$$

where $N_c = 3$, and Λ_0 denotes the condensation scale which is equal to Λ_h in the present case.⁴ The scalar potential of ϕ is not affected significantly by W_{np} as long as $H_{\text{inf}} \phi_0^2 \gg \Lambda_0^3$, which is satisfied for the parameters of our interest. Also, $f_a > \Lambda_0$ is needed in order not to destabilize the saxion, as will be shown below. Then the superpotential $W = \Delta W + W_{\text{np}}$ leads to the axion potential,

$$V(a) = 2N_f \sqrt{c-1} H_{\text{inf}} \Lambda_0^3 \cos\left(\frac{a}{N_c f_a} + \theta_0\right), \quad (7)$$

with the axion mass,

$$m_a^2 = \frac{\hat{c} H_{\text{inf}} \Lambda_0^3}{f_a^2}, \quad (8)$$

where $\hat{c} \equiv 4\sqrt{c-1} N_f / N_c^2$. The axion acquires a mass given by (8) as a result of the gluino condensation and SUSY breaking, as long as

$$H_{\text{inf}} < \Lambda_0 < f_a. \quad (9)$$

² If H_u and H_d are charged under the PQ symmetry, the effective axion decay constant during inflation is given by ϕ_0 . The isocurvature perturbation constraint can be similarly relaxed, if one introduces a singlet which gives a mass to additional quarks.

³ If the phase of ϕ is lighter than H_{inf} , it acquires quantum fluctuations, giving rise to the axion isocurvature perturbations with an amplitude suppressed by a factor of f_a/ϕ_0 compared to the conventional scenario.

⁴ The Higgs F -term $F_\phi/\phi \sim H_{\text{inf}}$ induces a gluino mass through gauge mediation. Thus H_{inf} should be smaller than Λ_0 in order for the gluino not to decouple at a scale above Λ_0 .

Note that a constant term in the superpotential $W \supset m_{3/2} M_{Pl}^2$ also contributes to the axion mass, which will be important when $m_{3/2} > H_{\text{inf}}$.

On the other hand, if there are n_f quarks lighter than Λ_h , they form a meson field stabilized by balancing the Affleck-Dine-Seiberg potential [20] and the SUSY mass term m . In practice, this is the case for $\Lambda_h \gtrsim 10^{11}$ GeV, where the up and down quarks are lighter than Λ_h . Integrating out the meson fields, we are left with the gluino condensation (6) with $\Lambda_0^{3N_c} = \Lambda_h^{3N_c - n_f} \det(m)$. Using Λ_0 defined this way, the axion mass formula (8) holds in this case.

The higher QCD scale and the heavy axion mass during inflation provide us with a simple way to suppress the axion CDM isocurvature perturbations. The axion mass is heavier than the inflation scale, i.e., $m_a > H_{\text{inf}}$, if

$$H_{\text{inf}} < 10^{12} \text{GeV} \hat{c} \left(\frac{\Lambda_0}{f_a} \right)^3 \left(\frac{f_a}{10^{12} \text{GeV}} \right). \quad (10)$$

Then the axion will be stabilized at one of the minima of the potential (7), and no sizable quantum fluctuations are generated at superhorizon scales. From (2) and (10), one can see that the isocurvature constraint on the inflation scale can be significantly relaxed compared to the conventional scenario.

So far we have assumed that f_a remains the same during and after inflation. This is the case if the saxion stabilization is not significantly modified by the strong QCD interactions, which generically requires $\Lambda_0 < f_a$. To see this, consider a model where the PQ symmetry is broken by two PQ fields S_1 and S_2 with the superpotential $W = \Sigma(S_1 S_2 - f_0^2)$ where Σ is a PQ singlet. For $\Lambda_0 < f_0$, the minimum will lie along the F -flat direction $S_1 S_2 = f_0^2$ while giving $f_a \sim f_0$, if S_1 and S_2 have soft scalar masses of similar size and couple to PQ charged quarks. On the other hand, the saxion potential is significantly affected for $\Lambda_0 > f_0$, as the non-perturbative superpotential gives the dominant contribution to the saxion potential. Note that, depending on the axion model, f_a may have a value different from the present one, even if $\Lambda_0 < f_a$.

Fig. 1 shows the upper bound on the inflation scale allowed by the constraint $m_a > H_{\text{inf}}$, taking $\hat{c} = 1$ and Λ_0 as a free parameter. We will consider an explicit model to realize a large




FIG. 1: The axion CDM isocurvature perturbation is suppressed by the heavy axion mass in the white region. See the text for explanation.

value of Λ_0 later. The axion CDM isocurvature perturbation is suppressed by the heavy axion mass in the white region. In the upper left shaded (gray) region, the saxion stabilization is significantly modified by the QCD interactions as $\Lambda_0 > f_a$. The thick (thin) solid line is the upper bound on H_{inf} for $\Lambda_0 = f_a$ ($\Lambda_0 = 0.1 f_a$). The dot-dashed (blue) lines represent the bound on H_{inf} for $\Lambda_0 = 10^8, 10^{10}, 10^{12}, 10^{14}$ GeV, from left to right. The upper right (yellow) region is excluded by the perturbativity limit of the gauge interactions up to the GUT scale (16). The conventional isocurvature bound (2) is shown by the dashed red line, where the anharmonic effect [7] is taken into account.

Let us here comment on the axion relic abundance from the misalignment mechanism. The axion is located at the minimum of the potential (7) after the inflation ends. Hence the axion dark matter abundance today reads

$$\Omega_a h^2 \simeq 0.2 \theta_i^2 \left(\frac{f_a}{10^{12} \text{GeV}} \right)^{1.184}, \quad (11)$$

with the initial misalignment angle given by

$$\theta_i = \theta_0 - \theta_{\text{QCD}}, \quad (12)$$

where θ_{QCD} denotes the QCD angle. For $f_a \gg 10^{12}$ GeV, $|\theta_i| \ll 1$ is needed to avoid overclosure of the Universe. In other words, the effect of the

stronger QCD is to suppress the axion quantum fluctuations, not the abundance.

A high QCD scale can be realized easily if there are extra colored particles that become heavier during inflation. Taking this into account, we consider the superpotential

$$W = \left(M_\Psi + \frac{\phi^2}{M'} \right) \Psi \Psi^c, \quad (13)$$

for N_Ψ pairs of $\Psi + \Psi^c$ that belong to $\mathbf{5} + \bar{\mathbf{5}}$ of SU(5) but do not carry PQ charges. The SM charged Ψ and Ψ^c are fixed at the origin if $M_\Psi + \phi^2/M'$ is larger than $m_{3/2}$ and H_{inf} . For ϕ fixed around the GUT scale, $\Psi + \Psi^c$ become heavy, and therefore the QCD confines at a larger scale than in the MSSM. Such effect is maximized when the effective cut-off scale M' is around M_{GUT} . We find that the QCD scale can be as high as,

$$\Lambda_h \simeq 1.3 \times 10^7 \text{GeV} \left(\frac{M_{\text{GUT}}}{M_\Psi} \right)^{N_\Psi/9}, \quad (14)$$

for $\phi_0 \sim M' \sim M_{\text{GUT}}$. Here we have taken into account the hierarchy in the SM quark masses, and used that the universal gauge coupling constant at the GUT scale is $g_{\text{GUT}}^2 \simeq 0.5$ in the MSSM.

It is also important to note that $\Psi + \Psi^c$ increase the axion mass since they decouple while giving a contribution to the coefficient of $\ln \phi$ in f_h in the effective theory. It is straightforward to see that $\hat{c}$ is enhanced by a factor $1 + 2N_\Psi/N_f$, and therefore the constraint on H_{inf} is further relaxed.

On the other hand, in the axion models where the PQ symmetry is linearly realized, there should exist PQ charged quarks in order for S to couple to the QCD anomaly. Let us introduce N_Φ pairs of $\Phi + \Phi^c$ transforming as $\mathbf{5} + \bar{\mathbf{5}}$ under SU(5), and consider the superpotential term $S\Phi\Phi^c$. Then $\Phi + \Phi^c$ obtain large masses $M_\Phi \sim f_a$. The gauge coupling unification is maintained in the presence of additional $\mathbf{5} + \bar{\mathbf{5}}$ matter fields, but the gauge coupling at the GUT scale is larger than the MSSM value. In order for the gauge interactions to remain perturbative up to M_{GUT} , we need

$$N_\Psi < \frac{151.6 - N_\Phi \ln(M_{\text{GUT}}/M_\Phi)}{\ln(M_{\text{GUT}}/M_\Psi)}. \quad (15)$$

The above is combined with the relation (14) to put the upper bound on the QCD scale,

$$\Lambda_h < 2.7 \times 10^{14} \text{GeV} \left(\frac{M_\Phi}{M_{\text{GUT}}} \right)^{N_\Phi/9}, \quad (16)$$




FIG. 2: Condensate scale Λ_0 for ϕ fixed at M_{GUT} in the presence of extra $\mathbf{5} + \bar{\mathbf{5}}$ matter fields. See the text for explanation.

which is insensitive to M_Ψ . Thus the QCD scale can be high enough to achieve $m_a > H_{\text{inf}}$ in the early Universe. Note that Λ_0 is slightly smaller than Λ_h for $\Lambda_h \gtrsim 10^{-5} M_{\text{GUT}}$, due to the hierarchy of the SM quark masses.

Fig. 2 shows how much Λ_0 can be raised for $\phi_0 = M_{\text{GUT}}$ during inflation, under the requirement of the perturbativity of gauge couplings up to the GUT scale. The solid (blue) lines represent Λ_0 for $N_\Psi = 3, 6, 10$ from bottom to top, while the dashed one is for the MSSM without extra matter fields. In the shaded region above the dot-dashed black line, the perturbativity of gauge couplings breaks down below M_{GUT} . This bound becomes severer in the presence of PQ charged fields $\Phi + \Phi^c$: the upper (lower) solid-black line is the bound for the case with $N_\Phi = 1$ and $M_\Phi = 10^{14}$ GeV ($M_\Phi = 10^{12}$ GeV).

The axion mass depends on c , the coupling of ϕ with the inflaton. If $c \gtrsim \mathcal{O}(10)$, the constraint on the Hubble parameter can be relaxed further, making the chaotic inflation consistent with the axion dark matter.

So far we have focused on the case of $m_a > H_{\text{inf}}$ during inflation. If this is not satisfied, the axion acquires quantum fluctuations. Note however that the axion may remain heavy for a while even after inflation. This is especially the case if the inflaton

decays into hidden particles, not the SM particles. Then, if the axion starts to oscillate and decays before ϕ starts to oscillate, the isocurvature constraint can be avoided.

Lastly let us mention the cosmological evolution after inflation. The Universe will be reheated by the decay of either the inflaton or ϕ . In the latter case, the preheating as well as subsequent dissipation effects will be important if ϕ passes through the origin during oscillations [21, 22].

In this letter we have proposed a novel mechanism to suppress the axion CDM isocurvature perturbations. The point is that, if QCD becomes strong at an intermediate or high scale during inflation, the axion can be so heavy that no sizable quantum fluctuations at superhorizon scales are produced.

Acknowledgment

This work was supported by Grant-in-Aid for Scientific Research (C) (No. 23540283) [KSJ], Scientific Research on Innovative Areas (No.24111702 [FT], No. 21111006 [FT], and No.23104008 [KSJ and FT]), Scientific Research (A) (No. 22244030 and No.21244033) [FT], and JSPS Grant-in-Aid for Young Scientists (B) (No. 24740135) [FT].

-
- [1] R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38**, 1440 (1977); Phys. Rev. D **16**, 1791 (1977).
 - [2] For a review, see J. E. Kim, Phys. Rept. **150**, 1 (1987); H. Y. Cheng, Phys. Rept. **158**, 1

- (1988); J. E. Kim and G. Carosi, Rev. Mod. Phys. **82**, 557 (2010); A. Ringwald, Phys. Dark Univ. **1** (2012) 116; M. Kawasaki and K. Nakayama, arXiv:1301.1123 [hep-ph].
- [3] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1303.5082 [astro-ph.CO].
- [4] D. Seckel and M. S. Turner, Phys. Rev. D **32**, 3178 (1985).
- [5] D. H. Lyth, Phys. Lett. B **236**, 408 (1990).
- [6] D. H. Lyth, Phys. Rev. D **45**, 3394 (1992).
- [7] T. Kobayashi, R. Kurematsu and F. Takahashi, arXiv:1304.0922 [hep-ph].
- [8] A. D. Linde, Phys. Lett. B **129**, 177 (1983).
- [9] M. Kawasaki, M. Yamaguchi and T. Yanagida, Phys. Rev. Lett. **85**, 3572 (2000).
- [10] R. Kallosh and A. Linde, JCAP **1011**, 011 (2010).
- [11] K. Nakayama, F. Takahashi and T. T. Yanagida, arXiv:1303.7315 [hep-ph].
- [12] A. D. Linde, Phys. Lett. B **259**, 38 (1991).
- [13] S. Folkerts, C. Germani and J. Redondo, arXiv:1304.7270 [hep-ph].
- [14] G. R. Dvali, hep-ph/9505253.
- [15] T. Banks and M. Dine, Nucl. Phys. B **505**, 445 (1997).
- [16] K. Choi, H. B. Kim and J. E. Kim, Nucl. Phys. B **490**, 349 (1997).
- [17] I. Affleck and M. Dine, Nucl. Phys. B **249**, 361 (1985).
- [18] F. L. Bezrukov and M. Shaposhnikov, Phys. Lett. B **659**, 703 (2008).
- [19] K. Nakayama and F. Takahashi, JCAP **1211**, 007 (2012).
- [20] I. Affleck, M. Dine and N. Seiberg, Nucl. Phys. B **241**, 493 (1984).
- [21] J. Garcia-Bellido, D. G. Figueroa and J. Rubio, Phys. Rev. D **79**, 063531 (2009).
- [22] K. Mukaida and K. Nakayama, JCAP **1303**, 002 (2013).